

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

170 Naturalist's-Meteorological-Celestial Reports. [Aug.

NATURALIST'S REPORT.

	From the 20th July, to the 20th August, 1812.
July	21, Entire leaved Virgin's Bower (Clematis integrifolia), flowering.
•	24, Scarlet Lychnis (Lychnis Chalcedonica), and Nettle leaved Bell-flower
	(Campanula Trachelium), flowering.
	25, Spear Thistle (Carduus Lanceolatus), flowering.
	26, Common Loosestrife (Lysimachia Vulgaris), and Shrubby broad-leaved
	Hawkweed (Hieracium sabaudum), flowering.
	27, Glutinous Acacia (Robinia Glutinosa), flowering.
	30, Common Besom Heath (Calluna Vulgaris), and narrowest-leaved Wil-
	low Herb (Epilobium Angustissimum).
August	4, Marsh Lousewort (Pedicularis Palustris), Devil's Bit (Scabiosa Succissa),
•	flowering.
	7, Woolly leaved Spiræa (Spiræa Tomentosa), flowering.
	8, Wandering Heath (Erica vagans), and Upright Loosestrife (Lysimachia Stricta), flowering.
	4, Scarlet Turk's-Cap Lily (Lilium Chalcedonicum), flowering.
	, Willow leaved Loosestrife (Lysimachia Ephemerum), and Great flowered
	Lavatera (Lavatera Thuringiaca), flowering.
	20, Tyger Peacock flower (Tigridia Pavonia), after surviving through last
	winter in the open ground, in flower.

METEOROLOGICAL REPORT.

	From 20th July, to 20th August, 1812.
July	21,Fine light showers in the evening.
- 7	22,Some very light showers.
	23,Showery.
	24,Wet.
	25-28,Showery.
	29—31,Fine.
Anguist	1,Showery, wet night.
****	2,Cloudy morning; tine day,
	3,Fine day; showery evening.
	413,Fine.
	14, Very fine day; rain at night.
	15,Showery.
	16,Heavy Showers in the evening.
	17,Dry.
	18,Dry day; rain at night.
	19,Showery; wet, stormy night.
ard . 2	20,Fine day; wet night.
	ighest state of the Barometer was on the 13th of August, when it stood at
	the lowest 29-4; the rest of the time it was very variable.
	hermometer on the 16th of August stood as high as 84°, at 8, A.M.; 18th
19th, an	d 20th, as high as 63°; on the 21st, 27th, and 28th of July, and 8th of August

at 55%.

The wind during this period has been very variable; it was observed 4 times N.—7 S.—9 W.—5 N.W.—11 S.W.—1 E.—13 N.E.—7 S.E.

CELESTIAL PHENOMENA.

FOR SEPTEMBER, 1812.

We have noticed the propriety of every one who remarks a peculiar appearance in the heavens sending an account of it to some one versed in astronomy, as every obser-